

1-9 HW Answer Key

2. $(2x+3)(x-2)(x+2)$

3. $x^2(3x-2)(3x-2)$

4. $\{1/2\}$

5. $\{3/2, -4\}$

6. $\{-2, \pm 1\}$

7. $\{0, 5, 2\}$

8. $\{0, \pm\sqrt{3}, \pm\sqrt{5}\}$

9. $\{\pm 3\}$

10. $\{\pm 2, \pm 5\}$

11. $(2x-5)(4x^2+10x+25)$

1. Roots and zeros are similar in that they both refer to the same point graphically where the function or related function crosses the x-axis. They are different in that roots are algebraic solutions to a given equation or an equation formed by setting the function equal to zero. Zeros are where the function crosses the x-axis (x-intercepts) or where the y value is zero.

Sep 20-4:40 PM

1. Explain what is similar and what is different between roots and zeros.

Roots and zeros are similar because they both refer to the same point graphically where the function or related function crosses the x-axis. They are different because roots are algebraic solutions to a given equation or an equation formed by setting the function equal to zero. Zeros are where the function crosses the x-axis (x-intercepts) or where the y value is zero.

$$2. 2x^3 + 3x^2 - 8x - 12$$

$$= x^2(2x+3) - 4(2x+3)$$

$$= (2x+3)(x^2-4)$$

$$= (2x+3)(x+2)(x-2)$$

$$3. 9x^4 - 12x^3 + 4x^2$$

$$= x^2(9x^2 - 12x + 4)$$

$$= x^2(3x-2)^2$$

$$\text{or } x^2(3x-2)^2$$

For the following:

- a) Algebraically find the zeros of the function.
b) Check graphically on your calculator by sketching the function and labeling the x-intercepts/zeros.
c) Check algebraically by substituting your solutions into the equation.

4a. $f(x) = 4x^2 - 4x + 1$

$$4x^2 - 4x + 1 = 0$$

$$4x^2 - 2x - 2x + 1 = 0$$


$$2x(2x-1) - 1(2x-1) = 0$$

$$(2x-1)(2x-1) = 0$$

$$2x-1 = 0 \quad | \quad x = 1/2$$

$$2x-1 = 0 \quad | \quad x = 1/2$$

$$x = 1/2$$



c. Algebraic check

$$4\left(\frac{1}{2}\right)^2 - 4\left(\frac{1}{2}\right) + 1 = 0$$

$$4\left(\frac{1}{4}\right) - 2 + 1 = 0$$

$$1 - 2 + 1 = 0$$

$$0 = 0 \quad \checkmark$$

5a. $f(x) = 2x^2 + 5x - 12$

$$2x^2 + 5x - 12 = 0$$

$$2x^2 + 8x - 3x - 12 = 0$$


$$2x(x+4) - 3(x+4) = 0$$

$$(x+4)(2x-3) = 0$$

$$x+4 = 0 \quad | \quad 2x-3 = 0$$

$$x = -4 \quad | \quad x = 3/2$$

$$x = -4, 3/2$$



c. Algebraic check

$$2(-4)^2 + 5(-4) - 12 = 0$$

$$32 - 20 - 12 = 0$$

$$0 = 0 \quad \checkmark$$

$$2\left(\frac{3}{2}\right)^2 + 5\left(\frac{3}{2}\right) - 12 = 0$$

$$2\left(\frac{9}{4}\right) + \frac{15}{2} - 12 = 0$$

$$\frac{9}{2} + \frac{15}{2} - 12 = 0$$

$$\frac{24}{2} - 12 = 0$$

$$12 - 12 = 0$$

$$0 = 0 \quad \checkmark$$

Sep 1-4:30 PM

6a. $f(x) = x^3 + 2x^2 - x - 2$
 $x^3 + 2x^2 - x - 2 = 0$
 $x^2(x+2) - 1(x+2) = 0$
 $(x+2)(x^2-1) = 0$
 $(x+2)(x+1)(x-1) = 0$
 $x = -2, -1, 1$
 $\{-2, \pm 1\}$

b. Graphic check

c. Algebraic check
 $(-2)^3 + 2(-2)^2 - (-2) - 2 = 0$
 $-8 + 8 + 2 - 2 = 0$
 $0 = 0 \checkmark$
 $(1)^3 + 2(1)^2 - (1) - 2 = 0$
 $1 + 2 - 1 - 2 = 0$
 $0 = 0 \checkmark$
 $(-1)^3 + 2(-1)^2 - (-1) - 2 = 0$
 $-1 + 2 + 1 - 2 = 0$
 $0 = 0 \checkmark$

Find the zeros of the following functions by factoring. No checks.

7. $f(x) = 3x^6 - 21x^5 + 30x^4$
 $3x^4(x^2 - 7x + 10) = 0$
 $3x^4(x-5)(x-2) = 0$
 $3x^4 = 0 \mid x = 5 \mid x = 2$
 $x = 0$
 $\{0, 5, 2\}$

8. $f(m) = m^5 - 8m^3 + 15m$
 $m^5 - 8m^3 + 15m = 0$
 $m(m^4 - 8m^2 + 15) = 0$
 $m(m^2-3)(m^2-5) = 0$
 $m = 0 \mid m^2-3=0 \mid m^2-5=0$
 $m = 0 \mid m = \pm\sqrt{3} \mid m = \pm\sqrt{5}$
 $\{0, \pm\sqrt{3}, \pm\sqrt{5}\}$

Find the roots of the following equations by factoring. No checks.

9. $3x^2 = 27$
 $3x^2 - 27 = 0$
 $3(x^2 - 9) = 0$
 $3(x+3)(x-3) = 0$
 $x = -3 \mid x = 3$
 $\{\pm 3\}$

10. $x^4 + 100 = 29x^2$
 $x^4 - 29x^2 + 100 = 0$
 $(x^2 - 25)(x^2 - 4) = 0$
 $(x+5)(x-5)(x+2)(x-2) = 0$
 $x = -5 \mid x = 5 \mid x = -2 \mid x = 2$
 $\{\pm 5, \pm 2\}$

11. Factor and check: $8x^3 - 125$
 $8x^3 - 125 = (2x-5)(4x^2 + 10x + 25)$
 $a = 2x$
 $b = 5$
 $(a-b)(a^2 + ab + b^2) = 8x^3 - 125 \checkmark$

check $(2x-5)(4x^2 + 10x + 25)$
 $= 8x^3 + 20x^2 + 50x - 20x^2 - 50x - 125$
 $= 8x^3 - 125 \checkmark$

Sep 1-4:30 PM

QUIZ

Sep 17-8:50 PM

1-10: Simplify Square Roots And Find The Discriminant

No calculator use. Complete and get familiar with this table.

x	x^2	x	x^2
Integer	Square	Integer	Square
1	1	11	121
2	4	12	144
3	9	13	169
4	16	14	196
5	25	15	225
6	36	16	256
7	49	17	289
8	64	20	400
9	81	25	625
10	100	30	900

Aug 27-1:49 PM

$$\sqrt[n]{x}$$

$n \rightarrow$ index
 $x \rightarrow$ radicand
 $\sqrt{} \rightarrow$ radical symbol

$$\sqrt{a}$$

$$\sqrt[3]{8}$$

Properties (true for all indices)

$$\sqrt{ab} = \sqrt{a} \cdot \sqrt{b}$$

$$\begin{aligned}
 \sqrt{4 \cdot 9} &= \sqrt{4} \cdot \sqrt{9} \\
 &= 2 \cdot 3 \\
 &= 6
 \end{aligned}$$

$$\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}} \quad \checkmark$$

$$\sqrt{\frac{100}{25}} = \frac{\sqrt{100}}{\sqrt{25}}$$

$$\sqrt{4} = \frac{10}{5}$$

$$2 = 2$$

Aug 27-1:50 PM

SIMPLIFY SQUARE ROOTS

Write each of the following numbers in simplest radical form:

$$1. \sqrt{24} = \sqrt{4 \cdot 6} = 2\sqrt{6}$$

$$2. \sqrt{16} = 4$$

$$3. \frac{\sqrt{54}}{\sqrt{2}} = \sqrt{\frac{54}{2}} = \sqrt{27} = \sqrt{9 \cdot 3} = 3\sqrt{3}$$

$$4. \sqrt{\frac{1}{225}} = \frac{\sqrt{1}}{\sqrt{225}} = \frac{1}{15}$$

$$5. \sqrt{\frac{72}{24}} = \sqrt{3} = \sqrt{3}$$

$$6. \frac{2}{3}\sqrt{27} = \frac{2}{3} \cdot \sqrt{9 \cdot 3} = \frac{2}{3} \cdot \frac{3\sqrt{3}}{1} = 2\sqrt{3}$$

Aug 27-2:03 PM

$$7. \sqrt{21} \text{ can't simplify!}$$

$$\begin{array}{r} \sqrt{4} \sqrt{12} \\ 2 \sqrt{4} \sqrt{3} \\ 2 \cdot 2 \sqrt{3} \\ \underline{4\sqrt{3}} \end{array}$$

$$8. -2\sqrt{48} = -2\sqrt{16 \cdot 3} = -2(4)\sqrt{3} = -8\sqrt{3}$$

$$9. 5\sqrt{16} = 5 \cdot 4 = 20$$

$$10. \sqrt{\frac{81}{16}} = \frac{\sqrt{81}}{\sqrt{16}} = \frac{9}{4}$$

Aug 27-1:51 PM

FIND THE DISCRIMINANT

We know that standard form of a quadratic equation is $ax^2 + bx + c = 0$

The discriminant is a number formed from standard form of a quadratic equation and equal to $b^2 - 4ac$.

Find the discriminant.

$$11. \overset{a}{x^2} - \overset{b}{3x} + \overset{c}{2} = 0$$

$$\begin{aligned} b^2 - 4ac &= (-3)^2 - 4(1)(2) \\ &= 9 - 8 \\ &= \boxed{1} \end{aligned}$$

$$12. \overset{a}{3x^2} - \overset{b}{5x} + \overset{c}{1} = 0$$

$$\begin{aligned} b^2 - 4ac &= (-5)^2 - 4(3)(1) \\ &= 25 - 12 \\ &= \boxed{13} \end{aligned}$$

Aug 27-1:51 PM

$$13. 4x^2 - 4x = -1$$

$$4x^2 - 4x + 1 = 0$$

$$\begin{aligned} b^2 - 4ac &= (4)^2 - 4(4)(1) \\ &= 16 - 16 \\ &= \boxed{0} \end{aligned}$$

$$14. x^2 = 7$$

$$x^2 - 7 = 0$$

$$\begin{aligned} b^2 - 4ac &= 0^2 - 4(1)(-7) \\ &= \boxed{28} \end{aligned}$$

$$15. x^2 - 6x + 9 = 0$$

$$\begin{aligned} b^2 - 4ac &= (-6)^2 - 4(1)(9) \\ &= 36 - 36 \\ &= \boxed{0} \end{aligned}$$

$$16. 2x^2 + 4x = 1$$

$$2x^2 + 4x - 1 = 0$$

$$\begin{aligned} b^2 - 4ac &= 4^2 - 4(2)(-1) \\ &= 16 + 8 = \boxed{24} \end{aligned}$$

Aug 27-1:59 PM

Simplify the square root of each discriminant in the previous section.

11. $\sqrt{1} = 1$

12. $\sqrt{13}$

13. $\sqrt{0} = 0$

14. $\sqrt{28} = \sqrt{4}\sqrt{7} = 2\sqrt{7}$

15. $\sqrt{0} = 0$

16. $\sqrt{24} = \sqrt{4}\sqrt{6} = 2\sqrt{6}$

Aug 27-1:59 PM

Sep 18-4:44 PM